Microwave-assisted synthesis of 2-acetyl-5-arylthiophenes and 4-(5-arylthiophen-2-yl)thiazoles via Suzuki coupling in water

Kamal M. Dawood,^{*a,b} Manahil B. Elamin,^a and Ahmad M. Farag^a

^aDepartment of Chemistry, Faculty of Science, Cairo University, Giza 12613, Egypt

^bCurrent address: Chemistry Department, Faculty of Science, Kuwait University, P.O. Box 5969, Safat 13060, Kuwait

E-mail: dr_dawood@yahoo.com

DOI: <http://dx.doi.org/10.3998/ark.5550190.p009.018>

Abstract

2-Acetyl-5-bromothiophene and 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole, as deactivated bromide candidates, were prepared and used for Suzuki cross-coupling reactions with a number of aryl(hetaryl)boronic acids in water or DMF as solvents. The cross-coupling reactions were carried out under thermal heating as well as microwave irradiating conditions using a benzothiazole-based Pd(II)-precatalyst. Optimization of the catalytic reaction condition was also studied.

Keywords: Thiophene, thiazole, palladium catalysis, microwave, Suzuki coupling

Introduction

Thiophene derivatives are important class of heterocyclic compounds that are widely involved in many agrochemicals and pharmaceuticals.¹ They are employed in drug synthesis, for example *Gabitril* is an antiepilepsy drug² and *Canagliflozin* is a drug for the treatment of type 2 diabetes³ (Chart 1). In addition, thiophene-based molecules have shown numerous biological activities such as antitumor,⁴ analgesic,⁵ anti-inflammatory,⁶ and antibacterial against Gram-positive bacteria.⁷ Suzuki-Miyaura reactions are among the most powerful tools for construction of carbon-carbon bond in organic synthesis both in industry and academia.⁸⁻¹¹ These reactions have also been widely exploited in the synthesis of natural products,^{12,13} and the design of pharmaceuticals.¹⁴ The use of water as green solvent has been encouraged by the desire to create cleaner, safer, and more environmentally benign chemical processes over the past few decades.¹⁵⁻¹⁷ Microwave irradiation methodology has also been receiving much interest in accelerating organic synthesis.^{18,19} 2-Acetyl-5-arylthiophenes were synthesized in literature via different routes such as: (1) direct arylation of 2-acetylthiophene with bromobenzenes,²⁰ or with

arene diazonium chlorides;²¹ (2) reaction of β -chloro- β -arylacroleins with sodium sulfide and chloroacetone;²² (3) cross-coupling of 2-acetyl-5-halothiophene with potassium aryl(heteroaryl)trifluoroborate,²³ or with aryl(heteroaryl)boronic acids,²⁴⁻²⁷ or with sodium tetraphenylborate;²⁸ (4) addition of acetonitrile to 2-phenylthiophene.²⁹ Recently, we reported the utility of several Pd(II)-complexes in C-C cross coupling reactions in water under microwave irradiation conditions,³⁰⁻³⁹ in this work; Suzuki cross-coupling reactions between two bromothiophene substrates having either activating (2-acetyl) or deactivating (2-thiazol-4-yl) moieties with variety of arylboronic acids under thermal heating as well as microwave irradiation conditions are evaluated using the benzothiazole-oxime Pd(II)-complex **1**.

Chart 1. Thiophene containing drugs; *Gabitril* and *Canagliflozin*.

Results and Discussion

Suzuki coupling of 2-acetyl-5-bromothiophene (**2**) with arylboronic acids under thermal heating and microwave irradiation

The first bromide substrate; 2-acetyl-5-bromothiophene (**2**) was prepared by treatment of 2-acetylthiophene with *N*-bromosuccinimide in a mixture of glacial acetic acid and acetic anhydride according to the reported literature.^{40,41} Suzuki cross-coupling reactions of 2-acetyl-5-bromothiophene (**2**) with activated and deactivated aryl(heteroaryl)boronic acids **3a-e** were carried out under thermal heating as well as microwave irradiation conditions using Pd(II)-precatalyst **1** as shown in Table 1. The cross-coupling reactions yielded of the corresponding 5-aryl(heteroaryl)-2-acetylthiophenes **4-8** in almost full conversions with excellent isolated yields regardless of the heating mode. The reaction components molar ratios were as following; 1 mmol 2-acetyl-5-bromothiophene (**2**), 1.2 mmol arylboronic acids **3a-e**, 0.6 mmol tetrabutylammonium bromide (TBAB), 2 mmoles of KOH and 0.25 mol% of the precatalyst **1** in water (3 mL). The identity of the resulted coupling products was confirmed from their ¹H and ¹³C NMR and MS spectral data. Although, sulfur species are reported as potential catalyst poisons that cause a remarkable decrease in the catalytic activity of palladium catalysts,⁴² also the thiophene moiety was mentioned as poison of a perfluorinated Pd-complex,⁴³ the cross-coupling of 2-acetyl-5-bromothiophene (**2**) with 3-thienylboronic acid (**3e**) proceeded straightforward giving excellent yield of 2-acetyl-3',5'-bithiophene (**8**) after 10 hours of thermal heating and after 9 min of microwave irradiation (Table 1, entry 5).

Table 1. Suzuki coupling of 2-acetyl-5-bromothiophene (**2**) with arylboronic acids using Pd-complex **1** under thermal heating and microwave irradiation

Entry	Ar(Het)-B(OH) ₂	Products 4-8	Thermal heating time/h	yield% ^a	MW heating time/min	yield% ^a
1	3a	4	1	93	1	95
2	3b	5	5	97	7	92
3	3c	6	5	90	7	98
4	3d	7	7	89	9	93
5	3e	8	10	91	9	95

^a Reaction condition: Bromide/ boronic acid/ KOH/ TBAB /water (3 mL): 1/1.2/2/0.6, at 100 °C for recorded time. ^aAll values refer to the isolated yields.

Synthesis of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**)

The hitherto unreported, 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) was prepared, as shown in Scheme 1, using Hantzsch method involving the reaction of α -halocarbonyl compounds with thiourea or thioamides. Thus, treatment of 2-acetyl-5-bromothiophene (**2**) with bromine in the presence of AlCl₃ afforded 5-bromo-2-(bromoacetyl)thiophene (**9**) in 96% yield.⁴⁴ Reaction of **9** with thioacetamide in ethanol at reflux temperature yielded 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) in 75% yield (Scheme 1). The structure of **10** was confirmed from its elemental analyses and spectroscopic data (IR, MS, ¹H and ¹³C NMR) as mentioned in the experimental section.

Scheme 1. Preparation of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**).

Optimization of the catalytic condition of Suzuki coupling of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (10**) with phenylboronic acid under microwave irradiation**

Suzuki cross-coupling of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) having a deactivating heterocyclic ring (thiazole) was examined. At first, it was better to optimize the catalytic condition of the new bromide substrate **10** with phenylboronic acid. The efficiencies of different bases and solvents as well as the proper catalyst concentration in the coupling reaction between **10** and phenylboronic acid were investigated under microwave irradiation. As shown in Table 2, runs 1-3, water/TBAB/KOH catalytic reaction condition was applied under microwave irradiation at different concentrations of Pd-complex **1** (0.125, 0.25 and 0.5 mol%, respectively), unfortunately there was no coupled products under all these concentrations. Repeating the same reaction using a mixed solvent (DMF/water/TBAB/KOH) and 0.5 mol% of Pd-complex **1** resulted in only 20% yield of **11**. When DMF was used as a sole solvent and KOH as base with 0.25 mol% of Pd-complex **1**, 50% yield was obtained (run 5, Table 2). Replacing KOH by Cs_2CO_3 using Pd-catalyst **1** in 0.25 and 0.5 mol% concentrations, gave 50% and 72% yield, respectively (Table 2, runs 6,7). Raising the Pd-complex concentration into 1 mol% using DMF/ Cs_2CO_3 resulted in full conversion as detected by TLC, after 30 min of microwave irradiation, with 80% isolated yield of 2-methyl-4-(5-phenylthiophen-2-yl)-1,3-thiazole (**11**) (run 8, Table 2). To evaluate the effect of heating mode, the cross-coupling reaction was repeated using DMF/ Cs_2CO_3 and 1 mol% of the precatalyst **1** under thermal heating at 100 °C for a longer time (40 hours) where only traces of the product **11** were detected by TLC (run 9, Table 2).

Suzuki coupling of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (10**) with arylboronic acids **3a-e** under microwave irradiation.**

Next, the optimized catalytic condition above was generalized for coupling of further arylboronic acids **3b-e** with 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) using Pd(II)-complex **1** (1 mol%) in DMF/ Cs_2CO_3 . The reaction components molar ratios were as following; 1 mmol of bromide **10**, 1.2 mmol of arylboronic acids **3**, 2 mmoles of Cs_2CO_3 using 1 mol% of the complex **1** in DMF (3 mL). The coupling reaction was conducted in a closed-vessel under microwave irradiation to give the corresponding arylthiophenylthiazole derivatives **11-14** in moderate to good isolated yields as outlined in Table 3. Unfortunately, the coupling of the bromide **10** with the deactivated 3-thienylboronic acid **3e** to get the bi(thienyl)thiazole derivative **15** resulted in no cross-coupling reaction even after 45 min of microwave irradiation (entry 5, Table 3).

Table 2. Optimization of the catalytic condition of Suzuki coupling of 4-(5-bromothiophen-2-yl)-2-methylthiazole (**10**)

Run	Cat. mol%	Solvent	Base	Yield% ^a
1	0.125	H ₂ O/TBAB	KOH	0.0
2	0.25	H ₂ O/TBAB	KOH	0.0
3	0.5	H ₂ O/TBAB	KOH	0.0
4	0.5	H ₂ O/DMF(1:2)/TBAB	KOH	20
5	0.25	DMF	KOH	50
6	0.25	DMF	Cs ₂ CO ₃	50
7	0.5	DMF	Cs ₂ CO ₃	72
8	1.0	DMF	Cs ₂ CO ₃	80
9	1.0	DMF ^b	Cs ₂ CO ₃	traces

^aReaction condition: Bromide/ boronic acid/ base/ solvent (3 mL): 1/1.2/2, under microwave irradiation (160 °C, 250 Watt) for 30 min. ^bWhen repeated under thermal heating at 100 °C for 40 h only traces of product **11** were detected by TLC.

Table 3. Suzuki coupling of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) with arylboronic acids **3a-e** under microwave irradiation

Run	Ar(Het)-B(OH) ₂ 3a-e	Products 11-15	MW irradiation ^a time/min yield%	
1	3a	11	30	80 ^b
2	3b	12	35	60 ^b

Table 3 (continued)

Run	Ar(Het)-B(OH) ₂ 3a-e	Products 11-15	MW irradiation ^a time/min	yield%
3	3c	13	40	76 ^b
4	3d	14	45	63 ^b
5	3e	15	45	0

^aConditions: Bromide **10**: 1 mmol; arylboronic acids **3a-e**: 1.2 mmol; Cs₂CO₃: 2 mmol; DMF (3 mL); Pd-complex **1**: 1 mol%, microwave irradiation (160 °C, 250 Watt), all values refer to the isolated yields. ^bTraces of the starting materials were detected by TLC.

An alternate Pd-free synthesis of 2-methyl-4-(5-phenylthiophen-2-yl)-1,3-thiazole (**11**).

2-Methyl-4-(5-phenylthiophen-2-yl)-1,3-thiazole (**11**) that was synthesized above, was alternatively prepared by a palladium-free conventional chemical method. Thus, bromination of 2-acetyl-5-phenylthiophene (**4**) following the literature procedure⁴⁵ yielded 2-(bromoacetyl)-5-phenylthiophene (**16**) in 92% yield. The latter was then treated with thioacetamide in absolute ethanol at reflux to give 2-methyl-4-(5-phenylthiophen-2-yl)-1,3-thiazole (**11**) in 95% yield as shown in Scheme 2.

Scheme 2. Alternative Synthesis of 2-methyl-4-(5-phenylthiophen-2-yl)-1,3-thiazole (**11**).

Conclusions

Although both the coupling candidates 2-acetyl-5-bromothiophene (**2**) and 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) have a bromine atom attached to the thiophene moiety, the reactivity of the thiazole containing candidate **10** was sharply different from that of **2** in water under microwave irradiation condition. Optimization of the catalytic conditions led to exploring the suitable coupling condition for such deactivated thiazolyl bromides. The obtained findings will be a lead to further investigations for coupling reactions of related deactivated bromides.

Experimental Section

General. Melting points were determined in open glass capillaries with a Gallenkamp apparatus and are uncorrected. The infrared spectra were recorded in potassium bromide disks on a Pye-Unicam SP 3-300 and Shimadzu FTIR 8101 PC infrared spectrophotometer. NMR spectra were recorded with a Varian Mercury VXR-300 NMR spectrometer at 300 MHz (^1H NMR) and at 75.46 MHz (^{13}C NMR) using deuterated chloroform (CDCl_3) or dimethylsulphoxide ($\text{DMSO}-d_6$). Chemical shifts are quoted in δ and were related to that of the solvents. Mass spectra (EI) were obtained at 70 eV with a type Shimadzu GCMQP 1000 EX spectrometer. Microwave experiments were carried out using a CEM Discover LabmateTM microwave apparatus (300 W with ChemDriverTM Software). Synthesis of the Pd(II)-complex **1**,³⁶ 2-acetyl-5-bromothiophene (**2**),^{40,41} 5-bromo-2-(bromoacetyl)thiophene (**9**),⁴⁴ and 2-(bromoacetyl)-5-phenylthiophene (**16**)⁴⁵ were accomplished following the procedures reported in literature.

Synthesis of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (10). To a solution of 5-bromo-2-(bromoacetyl)thiophene (**9**) (9.9 g, 35 mmol) in absolute ethanol (20 mL), thioacetamide (2.62 g, 35 mmol) was added and the mixture was refluxed for 2 hr then left to cool to room temperature and finally poured onto cold aqueous ammonia (pH 10). The solid product so formed was filtered off, washed with water and dried. Recrystallization from ethanol afforded an off-white solid of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) (7.1 g, 78% yield). mp 76-78°C; IR (ν_{max} , cm^{-1}): 3085, 2930, 1610, 1440. ^1H NMR ($\text{DMSO}-d_6$) δ 2.66 (s, 3H, CH_3), 7.20 (d, 1H, J 3.9 Hz), 7.36 (d, 1H, J 3.9 Hz), 7.79 (s, 1H); ^{13}C NMR ($\text{DMSO}-d_6$) δ_c 18.6, 110.9, 112.7, 124.2, 131.3, 139.7, 147.3, 166.2; MS m/z (%) 261 ($\text{M}^+ + 2$, 42.1), 259 (M^+ , 53.7), 138 (58.1), 94 (66.8), 93 (52.9), 81 (47.3), 68 (100). Anal. Calcd for $\text{C}_8\text{H}_6\text{BrNS}_2$: C, 36.93; H, 2.32; N, 5.38; S, 24.65. Found: C, 36.56; H, 2.40; N, 5.32; S, 24.81%.

Suzuki-Miyaura coupling of 2-acetyl-5-bromothiophene (2) with arylboronic acids using Pd-complex 1 in water under thermal heating. General Procedure. A mixture of 2-acetyl-5-bromothiophene (**2**) (205 mg, 1 mmol) and the appropriate arylboronic acids **3a-e** (1.2 mmol), TBAB (194 mg, 0.6 mmol), palladium complex **1** (1.14 mg, 0.25 mol%), KOH (112 mg, 2 mmol), and water (3 mL) were heated under reflux for the appropriate reaction times as listed in Table 1 (monitored by TLC). The products were then extracted with ethylacetate (EtOAc) (3x20 mL). The combined organic extracts were dried over anhydrous MgSO_4 then filtered and the solvent was evaporated under reduced pressure. The residue was then subjected to separation *via* flash column chromatography with n-hexane/EtOAc (4:1) as an eluent to give the corresponding pure cross-coupled products **4-8**.

Suzuki-Miyaura coupling of 2-acetyl-5-bromothiophene (2) with arylboronic acids using Pd-complex 1 in water under microwave irradiation. General Procedure. 2-Acetyl-5-bromothiophene (**2**) (205 mg, 1 mmol) and arylboronic acids **3a-e** (1.2 mmol), TBAB (194 mg,

0.6 mmol), palladium complex **1** (1.2 mg, 0.25 mol%), KOH (112 mg, 2 mmol), and distilled water (3 mL) were mixed in a process glass vial. The vial was capped properly, and thereafter the mixture was heated under microwave irradiating conditions at 160°C and 250 Watt for the appropriate reaction time, as listed in Table 1. After the reaction was almost complete, the cross-coupled products were extracted and then subjected to separation *via* flash column chromatography as above to give the corresponding pure cross-coupled products **4-8**.

2-Acetyl-5-phenylthiophene (4). Yellow solid, mp 113-114 °C (Lit.²⁷ mp 109-110 °C); IR ($\nu_{\max}$, cm^{-1}): 1647, 1442, 1353, 1276, 756. ^1H NMR (CDCl_3) δ_{H} 2.41 (3H, s, COCH_3), 7.15 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 4.2 Hz, 1CH), 7.21-7.23 (3H_{arom}, m, 3CH), 7.45 (1H_{arom}, d, $^3J_{\text{HH}}$ 6.6 Hz, 1CH), 7.52 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 4.2 Hz, 1CH); MS (EI, 70 eV): m/z (%) 202 (M^+ , 68.2), 191 (37.6), 162 (11.5), 92 (56.2), 84 (35.3), 76 (54.5), 50 (100).

2-Acetyl-5-(4-chlorophenyl)thiophene (5). Greenish-yellow solid, mp 135-136 °C (Lit.²¹ mp 114-115 °C); IR ($\nu_{\max}$, cm^{-1}): 2962, 2873, 1647, 1442, 1276, 1099. ^1H NMR ($\text{DMSO}-d_6$) δ_{H} 2.55 (3H, s, COCH_3), 7.52 (2H_{arom}, d, $^3J_{\text{HH}}$ 7.7 Hz, 2CH), 7.66 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 3.9 Hz, 1CH), 7.79 (2H_{arom}, d, $^3J_{\text{HH}}$ 7.5 Hz, 2CH), 7.94 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 3.9 Hz, 1CH); MS (EI, 70 eV): m/z (%) 238 ($\text{M}^+ + 2$, 77.7), 236 (M^+ , 100), 193 (13.1), 151 (65.3), 149 (98.4), 114 (68.6), 78 (48.5).

2-Acetyl-5-(4-methoxyphenyl)thiophene (6). Pale yellow solid, mp 160-161 °C (Lit.²² mp 156-157 °C); IR ($\nu_{\max}$, cm^{-1}): 1643, 1442, 1249, 1022. ^1H NMR ($\text{DMSO}-d_6$): δ_{H} 2.49 (3H, s, COCH_3), 3.80 (3H, s, OCH_3), 7.02 (2H_{arom}, d, $^3J_{\text{HH}}$ 7.5 Hz, 2CH), 7.52 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 5.7 Hz, 1CH), 7.71 (2H_{arom}, d, $^3J_{\text{HH}}$ 7.5 Hz, 2CH), 7.91 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 5.7 Hz, 1CH); ^{13}C NMR ($\text{DMSO}-d_6$): δ_{C} 26.2, 55.3, 114.6, 123.6, 127.4, 135.1, 141.6, 160.0, 190; MS (EI, 70 eV): m/z (%) 232 (M^+ , 100), 202 (13.5), 189 (33.1), 145 (91.8), 114 (25.4), 102 (48.7), 93 (20.2), 74 (32.7).

2-Acetyl-5-(4-methylphenyl)thiophene (7). Off-white solid, mp 118-119 °C (Lit.²¹ mp 115-116 °C); IR ($\nu_{\max}$, cm^{-1}): 1647, 1442, 1280, 1029, 798. ^1H NMR ($\text{DMSO}-d_6$): δ_{H} 2.34 (3H, s, CH_3), 2.55 (3H, s, COCH_3), 7.26 (2H_{arom}, d, $^3J_{\text{HH}}$ 7.8 Hz, 2CH), 7.58 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 4.2 Hz, 1CH), 7.63 (2H_{arom}, d, $^3J_{\text{HH}}$ 7.8 Hz, 2CH), 7.91 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 4.2 Hz, 1CH); ^{13}C NMR ($\text{DMSO}-d_6$): δ_{C} 20.8, 26.3, 120.5, 124.3, 125.8, 129.9, 135.0, 138.8, 142.1, 151.6, 190.4; MS (EI, 70 eV): m/z (%) 216 (M^+ , 85.7), 201 (100), 171 (11.9), 128 (98.3), 114 (22.1), 100 (25.7), 85 (17.5).

2-Acetyl-3',5'-bithiophene (8). Pale green solid, mp 122-123 °C (Lit.²³ mp 119-120 °C); IR ($\nu_{\max}$, cm^{-1}): 3097, 1655, 1450, 1276, 1029. ^1H NMR ($\text{DMSO}-d_6$): δ_{H} 2.5 (3H, s, COCH_3), 7.52 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 4.2 Hz, 1CH), 7.67-7.70 (2H_{arom}, m, 2CH), 7.69 (1H_{thiophene}, d, $^3J_{\text{HH}}$ 4.2 Hz, 1CH), 8.0 (1H_{thiophene}, s, 1CH); ^{13}C NMR ($\text{DMSO}-d_6$): δ_{C} 26.2, 122.9, 124.8, 125.9, 127.9, 134.1, 134.9, 141.7, 146.3, 190.4; MS (EI, 70 eV): m/z (%) 208 (M^+ , 100), 165 (24.2), 121 (76.3), 96 (26.8), 82 (33.1), 68 (21.4).

Effect of base, solvent and Pd-complex **1 concentration on Suzuki coupling of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) with phenylboronic acid (**3a**) under microwave irradiation.**

A mixture of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) (266 mg, 1 mmol), phenylboronic acid (**3a**) (146 mg, 1.2 mmol), TBAB (194 mg, 0.6 mmol), palladium complex **1** (0.6 mg, 0.125 mol%) and KOH (112 mg, 2 mmol), in water (3 mL) was subjected to microwave

irradiation at 160 °C and 250 Watt for 30 minutes to give 4-(5-phenylthiophen-2-yl)-2-methyl-1,3-thiazole (**11**). The same experiment was repeated using palladium complex **1** in different concentration (0.25, 0.5, and 1 mol%) with respect to 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**11**) using different solvents and bases. The molar ratio of the reaction components were in all cases as follows; 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole, phenylboronic acid, TBAB, base, solvent 1 / 1.2 / 0.6 / 2 / 3 mL. The results of this study are outlined in Table 2.

Suzuki-Miyaura coupling of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (10**) with phenylboronic acid **3a** under thermal heating.**

A mixture of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) (266 mg, 1 mmol) and phenylboronic acid **3a** (146 mg, 1.2 mmol), TBAB (194 mg, 0.6 mmol), palladium complex **1** (4.6 mg, 1 mol%) and Cs₂CO₃ (652 mg, 2 mmol), in DMF (3 mL) were refluxed at 100 °C for 40 hours as listed in Table 2. The starting bromide 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) was almost completely recovered without reaction.

Suzuki-Miyaura cross-coupling of 4-(5-bromothiophen-2-yl)-2-methyl-1,3-thiazole (10**) with arylboronic acids **3a-e** under microwave irradiation.**

General Procedure: 4-(5-Bromothiophen-2-yl)-2-methyl-1,3-thiazole (**10**) (266 mg, 1 mmol) and the appropriate arylboronic acids **3a-e** (1.2 mmol), TBAB (194 mg, 0.6 mmol), palladium complex **1** (4.6 mg, 1 mol%), Cs₂CO₃ (652 mg, 2 mmol), and DMF (3 mL) were mixed in a process glass vial. The vial was capped properly, and thereafter the mixture was heated under microwave irradiating conditions at 160 °C and 250 Watt for the appropriate reaction time as listed in Table 3. After the reaction was almost complete the cross-coupled products were then extracted with EtOAc (3x20 mL). The combined organic extracts were dried over anhydrous MgSO₄ then filtered and the solvent was evaporated under reduced pressure. The residue was then subjected to separation via flash column chromatography with n-hexane/EtOAc (4:1) as an eluent to give the corresponding pure cross-coupled products **11-15**. The results of this study are outlined in Table 3.

4-(5-Phenylthiophen-2-yl)-2-methyl-1,3-thiazole (11**).** Yellow powder, mp 92-93 °C; IR ($\nu_{\max}$, cm⁻¹): 3093, 2933, 1610, 1440, 1238. ¹H NMR (CDCl₃): δ_{H} 2.78 (3H, s, CH₃) 7.21 (1H_{thiazole}, s, 1CH), 7.27-7.41 (5H_{arom+thiophene}, m, 5CH), 7.64 (2H_{arom}, d, ³J_{HH} 7.5 Hz, 2CH); ¹³C NMR (CDCl₃): δ_{C} 19.2, 110.8, 123.7, 124.9, 125.6, 126.2, 127.5, 128.9, 129.1, 133.4, 137.3, 143.8, 149.3, 166.3; MS (EI, 70 eV): m/z (%) 257 (M⁺, 85), 180 (40.3), 148 (22.6), 123 (26.5), 111 (53.9), 95 (43.6). Anal. Calcd for C₁₄H₁₁NS₂: C, 65.33; H, 4.31; N, 5.44; S, 24.92%. Found: C, 65.11; H, 3.89; N, 5.22; S, 25.07%.

4-(5-(4-Chlorophenyl)thiophen-2-yl)-2-methylthiazole (12**).** Brown solid, mp 118-120 °C; IR ($\nu_{\max}$, cm⁻¹): 3105, 1442, 1272, 1091, 794. ¹H NMR (CDCl₃): δ_{H} 2.58 (3H, s, CH₃), 7.23 (1H_{thiazole}, s, 1CH), 7.29 (1H_{thiophene}, d, ³J_{HH} 3.9 Hz, 1CH), 7.40 (2H_{arom}, d, ³J_{HH} 7.8 Hz, 2CH), 7.58 (2H_{arom}, d, ³J_{HH} 7.8 Hz, 2CH), 7.65 (1H_{thiophene}, d, ³J_{HH} 3.9 Hz, 1CH); ¹³C NMR (CDCl₃): δ_{C} 26.5, 124.1, 125.9, 162.7, 127.4, 129, 129.3, 131.8, 133.1, 133.3, 134.9, 143.4, 151.1, 190.4; MS (EI, 70 eV): m/z (%) 293 (M⁺+2, 23.2), 291 (M⁺, 51.9), 221 (100), 148 (69.1), 113 (33.6), 85

(21.3). Anal. Calcd for $C_{14}H_{10}ClNS_2$: C, 57.62; H, 3.45; N, 4.80; S, 21.98%. Found: C, 57.42; H, 3.64; N, 4.76; S, 21.68%.

4-(5-(4-Methoxyphenyl)thiophen-2-yl)-2-methylthiazole (13). Grey powder, mp 122-124 °C; IR ($\nu_{\max}$, cm^{-1}): 3097, 2931, 1604, 1434, 1249, 1180, 802. 1H NMR ($CDCl_3$): δ_H 2.77 (3H, s, CH_3), 3.84 (3H, s, CH_3O), 6.92 (2H_{arom}, d, $^3J_{HH}$ 8.1 Hz, 2CH), 7.15 (1H_{thiophene}, d, $^3J_{HH}$ 3.9 Hz, 1CH), 7.27 (s, 1H), 7.35 (1H_{thiophene}, d, $^3J_{HH}$ 3.6 Hz, 1CH), 7.56 (2H_{arom}, d, $^3J_{HH}$ 8.1 Hz, 2CH); ^{13}C NMR ($CDCl_3$): δ_c 19.2, 55.4, 110.5, 113.9, 114.3, 122.6, 124.7, 126.9, 127.7, 130.1, 136.5, 143.7, 149.5, 159.3, 166.1; MS (EI, 70 eV): m/z (%) 287 (M^+ , 31.5), 272 (19.1), 257 (34), 255 (100), 128 (37.2), 95 (41.6), 64 (88.2). Anal. Calcd for $C_{15}H_{13}NOS_2$: C, 62.69; H, 4.56; N, 4.87; S, 22.31%. Found: C, 61.63; H, 4.36; N, 4.51; S, 22.58%.

2-Methyl-4-(5-(4-tolyl)thiophen-2-yl)thiazole (14). Grey powder, mp 114-116 °C; IR ($\nu_{\max}$, cm^{-1}): 3093, 2916, 1496, 1438, 1284, 1161. 1H NMR ($CDCl_3$): δ_H 2.37 (3H, s, CH_3), 2.76 (3H, s, $COCH_3$), 7.18-7.26 (4H_{arom}, m, 4CH), 7.37 (1H_{thiophene}, d, $^3J_{HH}$ 3.9 Hz, 1CH), 7.53 (2H_{arom}, d, $^3J_{HH}$ 7.8 Hz, 2CH); ^{13}C NMR ($CDCl_3$): δ_c 19.2, 21.2, 110.6, 123.1, 124.6, 125.5, 126.1, 129.5, 129.7, 131.5, 136.9, 137.4, 143.9, 149.5, 166; MS (EI, 70 eV): m/z (%) 271 (M^+ , 20.6), 270 (100), 230 (26.8), 129 (10.4), 93 (5.2), 83 (5.6). Anal. Calcd for $C_{15}H_{13}NS_2$: C, 66.38; H, 4.83; N, 5.16; S, 23.63%. Found: C, 67.03; H, 4.62; N, 5.01; S, 23.84%.

Alternative synthesis of 2-methyl-4-(5-phenylthiophen-2-yl)-1,3-thiazole (11). To a solution of 2-(bromoacetyl)-5-phenylthiophene (**16**) (2.46 g, 9 mmol) in absolute ethanol (20 mL), thioacetamide (0.66 g, 9 mmol) was added. The reaction mixture was refluxed for two hours, left to cool to room temperature then poured onto cold aqueous ammonia solution (pH 10). The solid product that formed was filtered off, washed with water and dried. Recrystallization from ethanol afforded 2-methyl-4-(5-phenylthiophen-2-yl)-1,3-thiazole (**11**) in 2.2 g (95% yield). The obtained product was in complete agreement (mp and spectral data) with the product that obtained above *via* Suzuki coupling reaction (Table 3, run 1).

Supplementary Material

Supplementary material is available.

References

1. Swanston, J. *Thiophene* In *Ullmann's Encyclopedia of Industrial Chemistry* Wiley-VCH: Weinheim, 2006.
2. Stahl, S. *Stahl's Essential Psychopharmacology: Prescriber's Guide*. Cambridge University Press: New York, NY, 2009; pp 523.
3. Chao, E. C. *Drug Fut.* **2011**, 36, 351.

- <http://dx.doi.org/10.1358/dof.2011.036.05.1590789>
4. Russel, K. R.; Press, B. J.; Rampulla, A. R. *J. Med. Chem.* **1988**, *31*, 1786.
<http://dx.doi.org/10.1021/jm00117a019>
 5. Kaur, H.; Kumar, S.; Sing, I.; Saxena, K. K.; Kumar, A. *Dig. J. Nanomater. Bios.* **2010**, *5*, 67.
 6. Badar, S. M. I. *Turk. J. Chem.* **2001**, *35*, 131.
 7. Sigmundova, I.; Zahradnik, P.; Magdolen, P.; Bujdakova, H. *Arkivoc* **2008**, (viii), 183.
<http://dx.doi.org/10.3998/ark.5550190.0009.815>
 8. Alonso, F.; Beletskaya, I. P.; Yus, M. *Tetrahedron* **2008**, *64*, 3047.
<http://dx.doi.org/10.1016/j.tet.2007.12.036>
 9. Bellina, F.; Carpita, A.; Rossi, R. *Synthesis* **2004**, *15*, 2419.
 10. Miyaura, N. *J. Organomet. Chem.* **2002**, *653*, 54.
[http://dx.doi.org/10.1016/S0022-328X\(02\)01264-0](http://dx.doi.org/10.1016/S0022-328X(02)01264-0)
 11. Suzuki, A. *J. Organomet. Chem.* **2002**, *653*, 83.
[http://dx.doi.org/10.1016/S0022-328X\(02\)01269-X](http://dx.doi.org/10.1016/S0022-328X(02)01269-X)
 12. Kelly, T. R.; Fu, Y.; Sieglen, J. T.; de Silva, H. *Org. Lett.* **2000**, *2*, 2351.
<http://dx.doi.org/10.1021/ol006127a>
 13. Jeong, S. J.; Higuchi, R.; Miyamoto, T.; Ono, M.; Kuwano, M.; Mawatari, S. F. *J. Nat. Prod.* **2002**, *65*, 1344.
<http://dx.doi.org/10.1021/np010577+>
 14. Ankita, C.; Jha, K.; Sachin, K. *J. Adv. Sci.* **2012**, *3*, 3.
 15. Dupont, J.; Consorti, C. S.; Spencer, J. *Chem. Rev.* **2005**, *105*, 2527.
<http://dx.doi.org/10.1021/cr030681r>
 16. Farina, V. *Adv. Synth. Catal.* **2004**, *346*, 1553.
<http://dx.doi.org/10.1002/adsc.200404178>
 17. Li, C. J. *Chem. Rev.* **2005**, *105*, 3095.
<http://dx.doi.org/10.1021/cr030009u>
 18. Hoz, A.; Ortiz, A. D.; Moreno, A. *Chem. Soc. Rev.* **2005**, *34*, 164.
<http://dx.doi.org/10.1039/b411438h>
 19. Kappe, C. O.; Stadler, A. *Microwave Theory*, In: *Microwaves in Organic and Medicinal Chemistry*; Wiley-VCH: Weinheim, 2005; Chapter 2.
 20. Churruca, F.; Hernandez, S.; Perea, M.; SanMartin, R.; Dominguez, E. *Chem. Commun.* **2013**, *49*, 1413.
<http://dx.doi.org/10.1039/c2cc37905h>
 21. Matyichuk, V. S.; Obushak, N. D.; Lytvyn, R. Z.; Horak, Yu. I. *Chem. Heterocycl. Compds.* **2010**, *46*, 50.
<http://dx.doi.org/10.1007/s10593-010-0468-3>
 22. Herbivo, C.; Comel, A.; Kirsch, G.; Manuela, M.; Raposo, M. *Tetrahedron* **2009**, *65*, 2079.
<http://dx.doi.org/10.1016/j.tet.2008.12.078>

23. Molander, G. A.; Biolatto, B. *J. Org. Chem.* **2003**, *68*, 4302.
<http://dx.doi.org/10.1021/jo0342368>
24. Kuriyama, M.; Matsuo, S.; Shinozawa, M.; Onomura, O. *Org. Lett.* **2013**, *15*, 2716.
<http://dx.doi.org/10.1021/ol4010189>
25. Barrett, M.; Fuchi, N. Jpn. Kokai Tokkyo Koho, JP 2013180955 A 20130912.
26. Molander, G. A.; Trice, S. L. J.; Kennedy, S. M. *J. Org. Chem.* **2012**, *77*, 8678.
<http://dx.doi.org/10.1021/jo301642v>
27. Kotha, S.; Kashinath, D.; Lahiri, K.; Sunoj, R. B. *Eur. J. Org. Chem.* **2004**, 4003.
<http://dx.doi.org/10.1002/ejoc.200400257>
28. Bai, L. *Chin. Chem. Lett.* **2009**, *20*, 158.
<http://dx.doi.org/10.1016/j.cclet.2008.10.050>
29. Jiang, T.-S.; Wang, G.-W. *Adv. Synth. Catal.* **2014**, 356, 369.
<http://dx.doi.org/10.1002/adsc.201300843>
30. Dawood, K. M.; Kirschning, A. *Tetrahedron* **2005**, *61*, 12121.
<http://dx.doi.org/10.1016/j.tet.2005.07.113>
31. Solodenko, W.; Brochwitz, C.; Wartchow, R.; Hashem, Md. A.; Vaultier, M.; Dawood, K. M.; Kirschning, A. *Mol. Div.* **2005**, *9*, 333.
<http://dx.doi.org/10.1007/s11030-005-7457-y>
32. Dawood, K. M.; Solodenko, W.; Kirschning, A. *Arkivoc* **2007**, (v), 104.
33. Dawood, K. M. *Tetrahedron* **2007**, *63*, 9642.
<http://dx.doi.org/10.1016/j.tet.2007.07.029>
34. Dawood, K. M.; Fayed, M. S.; Elkhalea, M. M. *Arkivoc* **2009**, (xiii), 324.
35. Shaaban, M. R.; Darweesh, A. F.; Dawood, K. M.; Farag, A. F. *Arkivoc* **2010**, (x), 208.
<http://dx.doi.org/10.3998/ark.5550190.0011.a18>
36. Darweesh, A. F.; Shaaban, M. R.; Farag, A. F.; Metz, P.; Dawood, K. M. *Synthesis* **2010**, 3163.
37. Dawood, K. M.; El-Deftar, M. M. *Arkivoc* **2010**, (ix), 319.
<http://dx.doi.org/10.3998/ark.5550190.0011.930>
38. Dawood, K. M.; El-Deftar, M. M. *Synthesis* **2010**, 1030.
<http://dx.doi.org/10.1055/s-0029-1218662>
39. Dawood, K. M.; Farag, A. F.; El-Deftar, M. M. Gardinar, M.; Abdelaziz, H. A. *Arkivoc* **2013**, (iii), 210.
<http://dx.doi.org/10.3998/ark.5550190.0014.317>
40. Smirnov, V. A.; Lipkin, A. E.; *Chem. Heterocycl. Compds.* **1975**, *9*, 170.
<http://dx.doi.org/10.1007/BF00569154>
41. Arsenyan, P.; Paegle, E.; Belyakov, S. *Tetrahedron Lett.* **2010**, *51*, 205.
<http://dx.doi.org/10.1016/j.tetlet.2009.10.133>
42. Dunleavy, J. K. *Platinum Metals Rev.* **2006**, *50*, 110.
<http://dx.doi.org/10.1595/147106706X111456>

43. Tzschucke, C. C.; Markert, C.; Glatz, H.; Bannwarth, W. *Angew. Chem. Int. Ed.* **2002**, *41*, 4500.
[http://dx.doi.org/10.1002/1521-3773\(20021202\)41:23<4500::AID-ANIE4500>3.0.CO;2-N](http://dx.doi.org/10.1002/1521-3773(20021202)41:23<4500::AID-ANIE4500>3.0.CO;2-N)
44. Perez, D. I.; Conde, S.; Pérez, C.; Gil, C.; Simon, D.; Wandosell, F.; Moreno, F. J.; Gelpi, J. L.; Luque, F. J.; Martinez, A. *Bioorg. Med. Chem.* **2009**, *17*, 6914.
<http://dx.doi.org/10.1016/j.bmc.2009.08.042>
45. Bagli, J. F.; Mackay, W. D.; Ferdinandi, E.; Cayen, M. N.; Vavra, I.; Pugsley, T.; Lippmann, W. *J. Med. Chem.* **1976**, *19*, 876.
<http://dx.doi.org/10.1021/jm00229a004>