One-pot five-component reaction for synthesis of some novel bis-dihydroquinazolinone derivatives

Ali A. Mohammadi,* Salman Tahery, and Saber Askari

Chemistry & Chemical Engineering Research Center of Iran, P.O. Box 14335-186, Tehran, Iran

E-mail: aliamohammadi@ccerci.ac.ir

DOI: <http://dx.doi.org/10.3998/ark.5550190.p008.715>

Abstract

An efficient and expedient procedure for the synthesis of some novel 3-(2-(4-oxo-2-aryl-1,2-dihydroquinazolin-3(4*H*)-yl)ethyl)-2-aryl-2,3-dihydroquinazolin-4(1*H*)-one and 3-(4-(4-oxo-2-aryl-1,2-dihydroquinazolin-3(4*H*)-yl)phenyl)-2-aryl-2,3-dihydroquinazolin-4(1*H*)-one derivatives is described. The method involves the one-pot five-component condensation of two molecules of isatoic anhydride, two molecules aldehyde with and one molecule of diamine in the presence of a catalytic amount of $\text{KAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ (alum). It affords the corresponding product in high yield with very short reaction time (50-75 min).

Keywords: Bisquinazolinone, aldehyde, isatoic anhydride, diamine, alum

Introduction

2,3-Dihydroquinazoline-4(3*H*)-ones are important fused heterocycles due to their potential biological and pharmaceutical activities. They have been used for the analgesic,¹ antitumor,² diuretic,³ antidefibrillatory,⁴ antihistamine,⁵ vasodilating agent,⁶ tranquilizer⁷ properties. Several methods have been reported for the synthesis of 2,3-dihydroquinazolinones,⁸⁻¹⁷ using various catalysts such as stannous chloride,¹⁸ ionic liquids,¹⁹ cyanuric chloride,²⁰ and Sc(III)-indapybox.²¹

The synthesis of bis-quinazoline derivatives was reported in only a few papers.²²⁻²⁷ The synthesis of 3,3'-(hexane-1,6-diyl)bis(2-(4-bromophenyl)-2,3-dihydroquinazolin-4(1*H*)-one) was described through a condensation catalyzed by iodine in ionic liquids²⁸ in very long reaction time (6-12 h) and 3-(2-(4-oxo-2-aryl-1,2-dihydroquinazolin-3(4*H*)-yl)ethyl)-2-aryl-2,3-dihydroquinazolin-4(1*H*)-one was reported by Reddy, et. al. in 1993 *via* two step in AcOH.²⁹

For wide variety of aromatic aldehydes (electron withdrawing or electron donating), the desired products were obtained in very short reaction time (within 50-75 min), good to excellent yields and simple workup procedure. The solid products obtained were just filtered of the reaction mixture. Water (25 mL) was added to the resulting solid (for removal of alum), filtered and washed with hot ethanol to give purified product. All products were characterized by their IR, MS, ^1H NMR, and ^{13}C NMR spectral data.

Table 1. Synthesis of bisquinazolinone **4a-r** using alum as catalysts

Products 4	Diamines 3	R	Time (min)	Yield (%) ^a	Mp (°C)	Lit. Yield (Lit. Time)	Lit. Mp (°C)
a	a	H	70	91	297-9	88 (3h) ²⁹	291
b	a	4-Cl	55	93	281-3	74 (3h) ²⁹	255
c	a	4-Me	55	96	295-7	85 (3h) ²⁹	270
d	a	4-NO ₂	55	96	286-8 (dec)	61(3h) ²⁹	275
e	a	2,4-diCl	50	90	310-13 (dec)	-	
f	a	3-EtO,4-OH	70	88	262-4	-	
g	a	3-Cl	55	93	264-6	64 (3h) ²⁹	245
h	a	4-MeO	55	94	250-2	69 (3h) ²⁹	260
i	a	4-CO ₂ H	70	90	308-10	-	
j	a	3-MeO	55	94	237-9	-	
k	a	2,4-diMeO	60	88	247-9	-	
l	b	H	75	90	296-8	-	
m	b	4-Cl	70	91	238-40 (dec)	-	
n	b	4-Me	65	93	281-3	-	
o	b	4-NO ₂	65	92	291-3	-	
p	b	4-MeO	65	90	251-3	-	
q	c	4-Me	75	90	182-4	88 (6h) ²⁸	179-81
r	c	4-Cl	65	85	213-5	79 (6h) ²⁸	212-4

^aIsolated yields for purified compounds.

Conclusions

In conclusion, the present method indicates an operationally simple and efficient one-pot five-component reaction for the preparation of bis quinazolin-4(3*H*)-one using the inexpensive, non-toxic, and easily available $\text{KAl}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ (alum) catalyst. In addition, low cost, excellent yields of products, easy experimental work-up procedure, and short reaction time make this methodology a valid contribution to the existing processes for the synthesis of bisquinazolin-4(3*H*)-one.

Experimental Section

General. Melting points were obtained in open capillary tubes and were measured on an electrothermal 9200 apparatus and are uncorrected. Mass spectra were recorded on a Shimadzu QP 1100 BX mass spectrometer. IR spectra were recorded on KBr pellets on a Shimadzu IR-470 spectrophotometer. ^1H and ^{13}C NMR spectra were determined on a Bruker 300 DRX Avance instrument at 300 and 75MHz. Elemental analysis for C, H and N were performed using a Heraeus CHN rapid analyzer. All the reactions are monitored by thin layer chromatography (TLC) with UV light as detecting agent.

General Procedure for the synthesis of bis(1,2-dihydro quinazolinon-4(1*H*)-one) derivatives (4 a-r). A mixture of isatoic anhydride **1** (2 mmol), aldehyde **2** (2 mmol), diamine **3** (1 mmol), 0.15 g (0.3 mmol) alum, and 10 mL EtOH 96% in a 50 mL flask was stirred at reflux for the time period as indicated in table 1. After completion of the reaction (monitored by TLC, ethyl acetate /*n*-hexane, 4:1), the solid products obtained were just filtered off the reaction mixture. Water (25 mL) was added to the resulting solid (for separation of alum), and the resulting solid was separated by filtration. The crude product was washed with hot ethanol to afford the purified product.

3,3'-(Ethane-1,2-diyl)bis(2-(2,4-dichlorophenyl)-2,3-dihydroquinazolin-4(1*H*)-one) (4e). Colourless solid; Yield: 88%; mp 310-13 °C (dec); IR (KBr): ν_{max} 3248 (NH), 3030, 1629 (C=O), 1521 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 2.86-2.90 (m, 2H, CH_2), 3.89-3.93 (m, 1H, CH_2), 3.96-4.00 (m, 1H, CH_2), 6.20 (d, 1H, J 2.6Hz, CH), 6.31 (d, 1H, J 2.7Hz, CH), 6.65-6.70 (m, 4H, Ar-H), 7.18-7.27 (m, 7H, 2NH, 5Ar-H), 7.35-7.37 (m, 1H, Ar-H), 7.63-7.65 (m, 4H, Ar-H) ppm; ^{13}C NMR (DMSO- d_6) δ 43.09, 68.3, 68.5, 114.9, 115.4, 118.3, 125.3, 128.3, 128.6, 129.2, 130.3, 134.4, 134.9, 143.9, 146.4, 163.1 ppm; MS: m/z (%) 614; Anal. Calcd for $\text{C}_{30}\text{H}_{22}\text{Cl}_4\text{N}_4\text{O}_2$: C, 58.84; H, 3.62; N, 9.15; Found: C, 58.76; H, 3.53; N, 9.07%.

3,3'-(Ethane-1,2-diyl)bis(2-(3-ethoxy-4-hydroxyphenyl)-2,3-dihydroquinazolin-4(1*H*)-one) (4f). Colourless solid; Yield: 81%; mp 262-4 °C; IR (KBr): ν_{max} 3397 (OH), 3281 (NH), 2976, 2929, 1634 (C=O), 1514 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 1.26 (t, 6H, J 7.0Hz, CH_3), 2.85-2.89 (m, 2H, CH_2), 3.91 (q, 4H, J 7.0Hz, CH_2), 4.02-4.04 (m, 2H, CH_2), 5.69 (s, 2H, CH), 6.61-6.71 (m,

8H, Ar-H), 6.90 (s, 2H, Ar-H), 7.17-7.19 (m, 4H, 2NH, 2Ar-H), 7.61 (d, 2H, *J* 8.0Hz, Ar-H), 8.99 (s, 2H, OH) ppm; ^{13}C NMR (DMSO-*d*₆) δ 15.5, 42.8, 64.6, 71.4, 112.8, 115.0, 115.5, 116.0, 117.8, 119.6, 128.2, 132.2, 134.0, 147.4, 147.5, 147.9, 163.3 ppm; MS: *m/z* (%) 594; Anal. Calcd for C₃₄H₃₄N₄O₆: C, 68.67; H, 5.76; N, 9.42; Found: C, 68.61; H, 5.66; N, 9.35%.

4,4'-(3,3'-(Ethane-1,2-diyl)bis(4-oxo-1,2,3,4-tetrahydroquinazoline-3,2-diyl))dibenzoic acid (4i). Colourless solid; Yield: 80%; mp 308-10 °C; IR (KBr): ν_{max} 3414 (OH), 3325 (NH), 2894, 1709 (C=O), 1693 (C=O), 1621 (C=O), 1568 cm⁻¹; ^1H NMR (DMSO-*d*₆) δ 2.9-3.00 (m, 2H, CH₂), 4.05-4.18 (m, 2H, CH₂), 5.95(d, 2H, *J* 2.1Hz, CH), 6.61 (d, 2H, *J* 8.0Hz, Ar-H), 6.66 (t, 2H, *J* 7.6Hz, Ar-H), 7.19 (t, 2H, *J* 7.0Hz, Ar-H), 7.39 (d, 4H, *J* 8.3Hz, Ar-H), 7.43 (d, 2H, *J* 2.1Hz, NH), 7.62 (d, 2H, *J* 6.8Hz, Ar-H), 7.86 (d, 4H, *J* 8.2Hz, Ar-H), 12.94 (broad, 2H, CO₂H) ppm; ^{13}C NMR (DMSO-*d*₆) δ 43.4, 70.6, 115.2, 115.5, 118.2, 127.2, 128.3, 130.4, 131.7, 134.2, 146.3, 147.0, 163.3, 167.6 ppm; MS: *m/z* (%) 562; Anal. Calcd for C₃₂H₂₆N₄O₆: C, 68.32; H, 4.66; N, 9.96; Found: C, 68.23; H, 4.68; N, 9.88%.

3,3'-(Ethane-1,2-diyl)bis(2-(3-methoxyphenyl)-2,3-dihydroquinazolin-4(1H)-one) (4j). Colourless solid; Yield: 94%; mp 237-9 °C; IR (KBr): ν_{max} 3240 (NH), 3002, 2930, 2828, 1631 (C=O), 1609 (C=O), 1514 cm⁻¹; ^1H NMR (DMSO-*d*₆) δ 2.89-2.94 (m, 1H, CH₂), 2.97-3.03 (m, 1H, CH₂), 3.68 (s, 6H, 2CH₃), 3.99-4.06 (m, 1H, CH₂), 4.10-4.18 (m, 1H, CH₂), 5.84 (d, 1H, *J* 1.5Hz, CH), 5.89 (d, 1H, *J* 1.7Hz, CH), 6.64 (t, 2H, *J* 8.3Hz, Ar-H), 6.69 (s, 2H, Ar-H), 6.85-6.89 (m, 6H, Ar-H), 7.18-7.27 (m, 4H, Ar-H), 7.36 (s, 1H, NH), 7.37 (s, 1H, NH), 7.63-7.65 (m, 2H, Ar-H) ppm; ^{13}C NMR (DMSO-*d*₆) δ 43.2, 43.4, 55.8, 71.2, 71.5, 113.13, 113.18, 114.3, 114.4, 115.1, 115.4, 115.5, 118.0, 119.0, 128.2, 130.5, 130.6, 134.1, 143.3, 147.3 160.2, 163.3, 163.4 ppm; MS: *m/z* (%) 534; Anal. Calcd for C₃₂H₃₀N₄O₄: C, 71.89; H, 5.66; N, 10.48; Found: C, 71.81; H, 5.59; N, 10.42%.

3,3'-(Ethane-1,2-diyl)bis(2-(2,4-dimethoxyphenyl)-2,3-dihydroquinazolin-4(1H)-one) (4k). Colourless solid; Yield: 90%; mp 247-9 °C; IR (KBr): ν_{max} 3384 (NH), 3067, 2936, 2837, 1649 (C=O), 1610, 1497 cm⁻¹; ^1H NMR (DMSO-*d*₆) δ 2.78-2.82 (m, 2H, CH₂), 3.68 (s, 3H, OCH₃), 3.81 (s, 3H, OCH₃), 3.96-4.00 (m, 2H, CH₂), 6.02 (d, 2H, *J* 1.9Hz, CH), 6.35 (d, d, 2H, *J* 2.3Hz, *J* 8.5Hz, Ar-H), 6.57 (d, 2H, *J* 2.3Hz, Ar-H), 6.62 (t, 2H, *J* 7.1Hz, Ar-H), 6.65 (d, 2H, *J* 8.0Hz, Ar-H), 6.81 (d, 2H, *J* 1.2Hz, 2NH), 6.92 (d, 2H, *J* 8.4Hz, Ar-H), 7.16 (t, 2H, *J* 8.3Hz, Ar-H), 7.61 (d, 2H, *J* 7.7Hz, Ar-H) ppm; ^{13}C NMR (DMSO-*d*₆) δ 42.9, 56.0, 56.5, 66.3, 99.6, 105.2, 115.2, 117.6, 120.9, 121.2, 127.7, 128.1, 133.9, 141.6, 158.5, 161.5, 163.7 ppm; MS: *m/z* (%) 594; Anal. Calcd for C₃₄H₃₄N₄O₆: C, 68.67; H, 5.76; N, 9.42; Found: C, 68.58; H, 5.68; N, 9.34%.

3,3'-(1,4-Phenylene)bis(2-phenyl-2,3-dihydroquinazolin-4(1H)-one) (4l). Cream solid; Yield: 82%; mp 296-8 °C; IR (KBr): ν_{max} 3305 (NH), 1638 (C=O), 1612, 1512 cm⁻¹; ^1H NMR (DMSO-*d*₆) δ 6.26 (s, 2H, CH), 6.70-6.75 (m, 4H, Ar-H), 7.24 (s, 4H, Ar-H), 7.29-7.33 (m, 12H, Ar-H), 7.66-7.70 (m, 4H, 2NH, 2Ar-H) ppm; MS: *m/z* (%) 522; Anal. Calcd for C₃₄H₂₆N₄O₂: C, 78.14; H, 5.01; N, 10.72; Found: C, 78.07; H, 4.92; N, 10.64%. ^{13}C NMR (125 MHz, DMSO-*d*₆) δ : very low soluble in DMSO.

3,3'-(1,4-Phenylene)bis(2-(4-chlorophenyl)-2,3-dihydroquinazolin-4(1H)-one) (4m).

Colourless solid; Yield: 87%; mp 238-40 °C(dec); IR (KBr): $\nu_{\max}$ 3305 (NH), 1641 (C=O), 1512 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 6.29 (d, 2H, J 2.3Hz, CH), 6.72 (t, 2H, J 7.3Hz, Ar-H), 6.74 (d, 2H, J 8.1Hz, Ar-H), 7.22 (s, 4H, Ar-H), 7.23-7.29 (m, 2H, Ar-H), 7.34-7.73 (m, 8H, Ar-H), 7.67 (d, 2H, J 2.3Hz, NH), 7.07 (d, 2H, J 7.0Hz, Ar-H) ppm; ^{13}C NMR (DMSO- d_6) δ 72.5, 115.7, 116.1, 118.6, 127.0, 128.8, 129.2, 129.3, 133.7, 134.8, 139.0, 140.5, 147.1, 162.9 ppm; MS: m/z (%) 590; Anal. Calcd for $\text{C}_{34}\text{H}_{24}\text{Cl}_2\text{N}_4\text{O}_2$: C, 69.04; H, 4.09; N, 9.47; Found: C, 68.95; H, 3.99; N, 9.39%.

3,3'-(1,4-phenylene)bis(2-(p-tolyl)-2,3-dihydroquinazolin-4(1H)-one) (4n).

Colourless solid; Yield: 92%; mp 281-3 °C; IR (KBr): $\nu_{\max}$ 3308 (NH), 3022, 2926, 1642 (C=O), 1611, 1512 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 2.22 (s, 6H, 2CH₃), 6.20 (s, 2H, CH), 6.70-6.72 (m, 4H, Ar-H), 7.09-7.23 (m, 10H, Ar-H), 7.54-7.77 (m, 8H, 2NH, 6Ar-H) ppm; MS: m/z (%) 550; Anal. Calcd for $\text{C}_{36}\text{H}_{30}\text{Cl}_2\text{N}_4\text{O}_2$: C, 78.52; H, 5.49; N, 10.17; Found: C, 78.44; H, 5.40; N, 10.10%.

^{13}C NMR (125 MHz, DMSO- d_6) δ : very low soluble in DMSO.

3,3'-(1,4-Phenylene)bis(2-(4-nitrophenyl)-2,3-dihydroquinazolin-4(1H)-one) (4o).

Yellow solid; Yield: 90%; mp 291-3 °C; IR (KBr): $\nu_{\max}$ 3401 (NH), 3102, 3071, 1662 (C=O), 1614, 1513 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 6.48 (d, 2H, J 2.1Hz, CH), 6.73 (t, 2H, J 7.4Hz, Ar-H), 6.76 (d, 2H, J 8.1Hz, Ar-H), 7.28 (t, 2H, J 7.2Hz, Ar-H), 7.33 (s, 4H, Ar-H), 7.62 (d, 4H, J 8.6Hz, Ar-H), 7.73 (d, 2H, J 7.6Hz, Ar-H), 7.82 (d, 2H, J 2.1Hz, 2NH), 8.15 (d, 4H, J 8.6Hz, Ar-H) ppm; ^{13}C NMR (DMSO- d_6) δ 72.3, 115.9, 116.0, 118.9, 124.5, 127.0, 128.6, 128.9, 134.9, 139.0, 146.8, 148.2, 148.8, 162.8 ppm; MS: m/z (%) 612; Anal. Calcd for $\text{C}_{34}\text{H}_{24}\text{N}_6\text{O}_6$: C, 66.66; H, 3.95; N, 13.72; Found: C, 66.56; H, 3.87; N, 13.63%.

3,3'-(1,4-Phenylene)bis(2-(4-methoxyphenyl)-2,3-dihydroquinazolin-4(1H)-one) (4p).

Colourless solid; Yield: 88%; mp 251-3 °C; IR (KBr) : $\nu_{\max}$ 3307 (NH), 2948, 2925, 2833, 1638, 1613, 1512 cm^{-1} ; ^1H NMR (DMSO- d_6) δ 3.68 (s, 6H, CH₃), 6.19 (d, 2H, J 2.5Hz, CH), 6.68 (t, 2H, J 7.1Hz, Ar-H), 6.74 (d, 2H, J 8Hz, Ar-H), 6.85 (d, 4H, J 8.7, Ar-H), 7.21 (s, 4H, Ar-H), 7.23-7.27 (m, 6H, Ar-H), 7.58 (d, 2H, J 2.4Hz, 2NH), 7.71 (d, 2H, J 7.0Hz, Ar-H) ppm; ^{13}C NMR (DMSO- d_6) δ 55.9, 72.9, 114.5, 115.6, 116.1, 118.3, 126.9, 128.5, 128.8, 133.5, 134.6, 139.2, 147.4, 159.9, 163.1 ppm; MS: m/z (%) 582; Anal. Calcd for $\text{C}_{36}\text{H}_{30}\text{N}_4\text{O}_4$: C, 74.21; H, 5.19; N, 9.62; Found: C, 74.14; H, 5.10; N, 9.54%.

Acknowledgements

We gratefully acknowledge financial support from the Research Council of Iran National Science Foundation: INSF.

Supplementary Information

General experimental procedures, IR, ^1H and ^{13}C NMR, and MS data for compounds **4 a-r** are available as supplementary information.

References

1. Hour, M.-J.; Huang, L.-J.; Kuo, S.-C.; Xia, Y.; Bastow, K.; Nakanishi, Y.; Hamel, E.; Lee, K.-H., *J. Med. Chem.* **2000**, *43*, 4479.
<http://dx.doi.org/10.1021/jm000151c>
2. Birch, H. L.; Buckley, G. M.; Davies, N.; Dyke, H. J.; Frost, E. J.; Gilbert, P. J.; Hannah, D. R.; Haughan, A. F.; Madigan, M. J.; Morgan, T.; Pitt, W. R.; Ratcliffe, A. J.; Ray, N. C.; Richard, M. D.; Sharpe, A.; Taylor, A. J.; Whitworth, J. M.; Williams, S. C., *Bioorg. Med. Chem. Lett.* **2005**, *15*, 5335.
<http://dx.doi.org/10.1016/j.bmcl.2005.06.108>
3. Cohen, E.; Klarberg, B.; Vaughan, J. R., *J. Am. Chem. Soc.* **1959**, *81*, 5508.
<http://dx.doi.org/10.1021/ja01529a062>
4. Bonola, G.; Da Re, P.; Magistretti, M. J.; Massarani, E.; Setnikar, I., *J. Med. Chem.* **1968**, *11*, 1136.
<http://dx.doi.org/10.1021/jm00312a007>
5. Alagarsamy, V.; Solomon, V. R.; Murugan, M., *Bioorg. Med. Chem.* **2007**, *15*, 4009.
<http://dx.doi.org/10.1016/j.bmc.2007.04.001>
6. Levin, J. I.; Chan, P. S.; Bailey, T.; Katocs Jr, A. S.; Venkatesan, A. M., *Bioorg. Med. Chem. Lett.* **1994**, *4*, 1141.
[http://dx.doi.org/10.1016/S0960-894X\(01\)80244-4](http://dx.doi.org/10.1016/S0960-894X(01)80244-4)
7. Hirose, N.; Kuriyama, S.; Sohda, S., *Chem. Pharm. Bull.* **1973**, *21*, 1005.
<http://dx.doi.org/10.1248/cpb.21.1005>
8. Ahmad, S.; Ngu, K.; Miller, K. J.; Wu, G.; Hung, C. p.; Malmstrom, S.; Zhang, G.; O'Tanyi, E.; Keim, W. J.; Cullen, M. J.; Rohrbach, K. W.; Thomas, M.; Ung, T.; Qu, Q.; Gan, J.; Narayanan, R.; Pelleymounter, M. A.; Robl, J. A., *Bioorg. Med. Chem. Lett.* **2010**, *20*, 1128.
<http://dx.doi.org/10.1016/j.bmcl.2009.12.014>
9. Badiger, D. S.; Nidavani, R. B.; Hunoor, R. S.; Patil, B. R.; Vadavi, R. S.; Chandrashekhar, V. M.; Muchchandi, I. S.; Gudasi, K. B. *Appl. Organomet. Chem.* **2011**, *25*, 876.
<http://dx.doi.org/10.1002/aoc.1855>
10. Bothara, K. G.; Kadam, S. S.; Sai Shivram, V. *Indian Drugs* **1998**, *35*, 372.
11. Burgey, C. S.; Stump, C. A.; Nguyen, D. N.; Deng, J. Z.; Quigley, A. G.; Norton, B. R.; Bell, I. M.; Mosser, S. D.; Salvatore, C. A.; Rutledge, R. Z.; Kane, S. A.; Koblan, K. S.; Vacca, J. P.; Graham, S. L.; Williams, T. M. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 5052.
<http://dx.doi.org/10.1016/j.bmcl.2006.07.044>

12. Chinigo, G. M.; Paige, M.; Grindrod, S.; Hamel, E.; Dakshanamurthy, S.; Chruszcz, M.; Minor, W.; Brown, M. L. *J. Med. Chem.* **2008**, *51*, 4620.
<http://dx.doi.org/10.1021/jm800271c>
13. Christie, R. M.; Moss, S. *J. Chem. Soc., Perkin Trans. 1* **1985**, 2779.
<http://dx.doi.org/10.1039/p19850002779>
14. Dabiri, M.; Salehi, P.; Baghbanzadeh, M. *Monatsh. Chem.* **2007**, *138*, 1191.
<http://dx.doi.org/10.1007/s00706-007-0635-0>
15. Sharma, R.; Pandey, A. K.; Chauhan, P. M. S. *Synlett* **2012**, *23*, 2209.
<http://dx.doi.org/10.1055/s-0032-1317014>
16. Sashidhara, K. V.; Palnati, G. R.; Dodda, R. P.; Avula, S. R.; Swami, P. *Synlett* **2013**, *24*, 1795.
<http://dx.doi.org/10.1055/s-0033-1339466>
17. Rostamizadeh, S.; Amani, A. M.; Mahdavinia, G. H.; Sepehrian, H.; Ebrahimi, S. *Synthesis* **2010**, *2010*, 1356.
18. Yoo, C. L.; Fettingner, J. C.; Kurth, M. J. *J. Org. Chem.* **2005**, *70*, 6941.
<http://dx.doi.org/10.1021/jo050450f>
19. Wang, X.-S.; Yang, K.; Zhou, J.; Tu, S.-J. *J. Comb. Chem.* **2010**, *12*, 417.
<http://dx.doi.org/10.1021/jo050450f>
20. Sharma, M.; Pandey, S.; Chauhan, K.; Sharma, D.; Kumar, B.; Chauhan, P. M. S. *J. Org. Chem.* **2011**, *77*, 929.
<http://dx.doi.org/10.1021/jo2020856>
21. Prakash, M.; Kesavan, V. *Org. Lett.* **2012**, *14*, 1896.
<http://dx.doi.org/10.1021/ol300518m>
22. Moghimi, A.; Khanmiri, R. H.; Omrani, I.; Shaabani, A. *Tetrahedron Lett.* **2013**, *54*, 3956.
<http://dx.doi.org/10.1016/j.tetlet.2013.05.065>
23. Lu, L.; Zhang, M.-M.; Jiang, H.; Wang, X.-S., *Tetrahedron Lett.* **2013**, *54*, 757.
<http://dx.doi.org/10.1016/j.tetlet.2012.11.042>
24. Wang, X. S.; Yang, K.; Zhou, J.; Tu, S. J. *J. Comb. Chem.* **2010**, *12*, 417.
<http://dx.doi.org/10.1021/cc900174p>
25. El-Sharief, A. M. S.; Ammar, Y. A.; Zahran, M. A.; Sabet, H. K. *Phosphorus, Sulfur, and Silicon and the Related Elements* **2004**, *179*, 267.
<http://dx.doi.org/10.1021/cc900174p>
26. Yang, X.-H.; Wu, M.-H.; Sun, S.-F.; Xie, J.-L.; Ding, M.-W.; Xia, Q.-H. *J. Heterocycl. Chem.* **2008**, *45*, 1365.
<http://dx.doi.org/10.1002/jhet.5570450518>
27. Helmholz, F.; Schroeder, R.; Langer, P. *Synthesis* **2006**, *2006*, 2507.
28. Liu, Y.; Lu, L.; Zhou, Y.-J.; Wang, X.-S., *Res. Chem. Intermed.* in press, DOI 10.1007/s11164-013-1131-2.
<http://dx.doi.org/10.1007/s11164-013-1131-2>
29. Zachariah, L.; Reddy, M. S., *Indian J. Chem.* **1993**, *32B*, 826.

30. Azizian, J.; Mohammadi, A. A.; Karimi, N.; Mohammadizadeh, M. R.; Karimi, A. R., *Catal. Commun.* **2006**, 7, 752.
<http://dx.doi.org/10.1016/j.catcom.2006.01.026>
31. Mohammadi, A. A.; Dabiri, M.; Qaraat, H., *Tetrahedron* **2009**, 65, 3804.
<http://dx.doi.org/10.1016/j.tet.2009.02.037>
32. Mohammadi, A. A.; Sadat Hossini, S. S. *Chin. J. Chem.* **2011**, 29, 1982.
<http://dx.doi.org/10.1002/cjoc.201180344>
33. Dabiri, M.; Salehi, P.; Otokesh, S.; Baghbanzadeh, M.; Kozehgary, G.; Mohammadi, A. A. *Tetrahedron Lett.* **2005**, 46, 6123.
<http://dx.doi.org/10.1016/j.tetlet.2005.06.157>
34. Azizian, J.; Mohammadi, A. A.; Karimi, A. E.; Mohammadizadeh, M. R. *J. Org. Chem.* **2005**, 70, 350.
<http://dx.doi.org/10.1021/jo049138g>
35. Azizian, J.; Mohammadi, A. A.; Karimi, A. R.; Mohammadizadeh, M. R. *Appl. Catal., A* **2006**, 300, 85.
36. Mohammadi, A. A.; Mivechi, M.; Kefayati, H. *Monatsh. Chem.* **2008**, 139, 935.
<http://dx.doi.org/10.1007/s00706-008-0875-7>