Synthesis of 2,2'-bis(1-(2-aryl)-1*H*-pyrazol-4-yl)-3,3,3',3'-tetramethyl-3*H*,3'*H*-5,5'-biindoles and 2,7-bis(1-(3-aryl)-1*H*-pyrazol-4-yl)-1,1,6,6-tetramethyl-1,6-dihydroindolo[7,6-*g*]indoles

Ahmad Rashidi,^a Mehdi M. Baradarani,^{a*} and John A. Joule^b

^aDepartment of Chemistry, Faculty of Science, University of Urmia, Urmia 57153-165, Iran

^bThe School of Chemistry, The University of Manchester, M13 9PL, UK

E-mail: mmbaradarani@yahoo.com

Abstract

2,2'-(Biphenyl-4,4'-diyl)bis(hydrazinium) chloride and 2,2'-(naphthalene-1,5-diyl)bis(hydrazinium) chloride were converted *via* Fischer syntheses with 3-methylbutan-2-one into bisindolenines, 2,2',3,3,3',3'-hexamethyl-3*H*,3'*H*-5,5'-biindole **10** and 2,3,3,7,8,8-hexamethyl-3*H*,8*H*-indolo[7,6-*g*]indole **14**, respectively. Exposure of the bisindolenines to the Vilsmeier reagent produced tetraformyl compounds **11** and **15**, which reacted with hydrazine and arylhydrazines to give the corresponding pyrazoles **12** and **16** in excellent yields.

Keywords: Arylhydrazines, bisindolenine, Fischer reaction, hydrazine, pyrazole, Vilsmeier reagent

Introduction

We recently described the reaction of 2,3,3-trimethylindolenines (3*H*-indoles) **1** with the Vilsmeier reagent formed from dimethylformamide and phosphorus oxychloride to produce aminomethylene malondialdehydes **2** (Scheme 1).^{1,2} Additionally we showed that the pyridoindolenines 2,3,3-trimethyl-3*H*-pyrrolo[2,3-*f*]quinoline and 2,3,3-trimethyl-3*H*-pyrrolo[3,2-*h*]quinoline **4** behave similarly (Scheme 2).³

Scheme 1

The malondialdehydes **2,5** reacted with hydrazine or arylhydrazines to produce substituted pyrazoles **3,6**.^{1,3,4} We have now been able to show that the principles embodied in these transformations can be incorporated into bisindolenine systems, and thus have prepared several more complex pyrazoles.

Scheme 2

For the mechanism of formation of the aminomethylene malondialdehydes, we suggested that a small equilibrium concentration of an enamine tautomer **7** is successively C-substituted and thus, before hydrolysis during work-up, species **8** is present (Scheme 3). We propose that a comparable mechanism operates in the work described herein.

Scheme 3

Results and Discussion

Diazotisation of benzidine and naphthalene-1,5-diamine, then reduction of the diazonium salts with tin(II) chloride, produced the corresponding bis hydrazinium chlorides **9** and **13**.⁵ Reaction of 2,2'-(biphenyl-4,4'-diyl)bis(hydrazinium)chloride **9** with isopropyl methyl ketone in a Fischer reaction⁶ produced the bisindolenine 2,2',3,3',3',3'-hexamethyl-3*H*,3'*H*-5,5'-biindole **10** in good yield (Scheme 4).

Scheme 4

Similarly, 2,2'-(naphthalene-1,5-diyl)bis(hydrazinium) chloride **13** reacted with isopropyl methyl ketone in hot acetic acid to give 2,3,3',7,8,8-hexamethyl-3*H*,8*H*-indolo[7,6-*g*]indole⁷ **14** (Scheme 5), which was used previously in the synthesis of photochromic compounds.⁸

The structure of the bisindolenine **10** was evident from its molecular formula, the twelve-hydrogen singlet for the geminal methyl groups, at δ 1.33 ppm, and a singlet signal for the imine methyl groups resonating at δ 2.30 ppm.

Each of the bisindolenines **10**, **14** was now reacted with the Vilsmeier reagent in yields of 90% and 85%, respectively, and tetraformyl compounds **11**, **15** were obtained (Schemes 4 and 5). The structures of the aminomethylene malondialdehydes rests on the observation of two two-hydrogen singlets at δ 9.79 and δ 9.83 for **11** and δ 9.83 and δ 9.87 for **15** corresponding to aldehyde protons. Absorptions at 3136 cm⁻¹ and 3130 cm⁻¹ for **11** and **15**, respectively, were evidence for the presence of N-H bonds, further confirmed by ¹H NMR two-hydrogen signals for the N-hydrogens appearing at δ 13.64 (**11**) and δ 14.35 (**15**), respectively. As in our previous work,^{1,3} the aminomethylene malondialdehydes reacted with hydrazine and various arylhydrazines to give pyrazoles, with migration of the double bond to reform the imine unit (Schemes 4 and 5). For pyrazoles **12a-f**, the newly formed five-membered heterocyclic ring protons resonated in the range δ 8.35-9.37 and for the pyrazoles **16a-e** in the range δ 8.50-9.38.

Scheme 5

Conclusions

We have been able to show that the principles embodied in transformations of simple indolenines *via* Vilsmeier formylations can be incorporated into more complex bisindolenine systems and thus have prepared several pyrazoles in excellent yields.

Experimental Section

General. Melting points were recorded on a Philip Harris C4954718 apparatus. ^1H and ^{13}C NMR spectra were recorded on a Bruker Avance AQS 300 MHz spectrometer, at 300 MHz and 75 MHz respectively. Chemical shifts δ are in parts per million (ppm) measured in CDCl_3 and $\text{DMSO}-d_6$ as solvents and relative to TMS as the internal standard. Infrared spectra were recorded on a Thermo Nicolet-Nexus 670 FT-IR instrument. High resolution mass spectra were recorded on an Agilent Technology (HP), MS Model: 5973 Network Mass, selective Detector Ion source: Electron Impact (EI) 70 eV, ion source temperature: 230°C Analyzer: quadrupole, and relative abundances of fragments are quoted in parentheses after the m/z values.

2,2',3,3',3',3'-Hexamethyl-3*H*,3'*H*-5,5'-biindole (10). A mixture of 2,2'-(biphenyl-4,4'-diyl)bis(hydrazinium)chloride (8.29 g, 28.9 mmol) and isopropyl methyl ketone (6.52 mL, 60.60 mmol) was heated at reflux in acetic acid (50 mL) for 12 h and then cooled, diluted with water (50 mL), and neutralized with NaOH 2M, then extracted with ethyl acetate (4 × 100 mL). The organic layer was dried over Na₂SO₄. The solvent was evaporated and the resulting viscous oil crystallized from EtOH to give the 2,2',3,3',3',3'-hexamethyl-3*H*,3'*H*-5,5'-biindole **10**. (6.38 g, Yield 70%); mp 104-106 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2962, 2927, 1685, 1574, 1458, 1409; ¹H NMR (CDCl₃): δ 1.33 (s, 12H), 2.30 (s, 6H), 7.49 (s, 2H), 7.53 (d, *J* = 8.1 Hz, 2H), 7.59 (d, *J* = 8.1 Hz, 2H); ¹³C NMR (CDCl₃): δ 15.45, 23.18, 53.75, 119.95, 120.30, 126.81, 138.70, 146.32, 152.99, 188.21; MS (EI, 70 ev): *m/z* (%) 316 (M⁺, 100), 301 (32), 285 (10), 84 (82), 47 (20). Found: [M]⁺ 316.1939, C₂₂H₂₄N₂ requires [M]⁺ 316.1939.

2,2'-(3,3',3',3'-Tetramethyl-5,5'-biindoline-2,2'-diylidene)dimalonaldehyde (11). To *N,N*-dimethylformamide (2 mL) cooled in an ice bath was added dropwise phosphorus oxychloride (1.6 mL, 17.4 mmol) with stirring at below 5 °C. After this addition, a solution of **10** (2.9 mmol, 0.918 g) in DMF (2 mL) was added dropwise. The cooling bath was removed and the reaction mixture was stirred at 75 °C for 12 h. The resulting solution was added to ice-cooled water and made alkaline with NaOH (aq.) solution. The resulting precipitate was collected by filtration after 24 h, dried in air, recrystallized from ethanol, to give **11** (1.12 g, Yield 90%); mp 213-215 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 3136, 2967, 2926, 2765, 1677, 1601, 1520, 1473, 1202; ¹H NMR (CDCl₃): δ 1.82 (s, 12H), 7.50 (s, 2H), 7.55 (d, *J* = 8.1 Hz, 2H), 7.59 (d, *J* = 8.1 Hz, 2H), 9.79 (s, 2H), 9.83 (s, 2H), 13.64 (bs, 2H); ¹³C NMR (CDCl₃): δ 23.46, 51.49, 109.65, 112.96, 120.96, 127.40, 138.50, 138.95, 141.65, 179.37, 187.81, 192.59; MS (EI, 70 ev): *m/z* (%) 428 (M⁺, 15), 400 (13), 361 (52), 291 (24), 263 (38), 248 (36), 191 (30), 149 (38), 57 (100), 41 (51). Found: [M]⁺ 428.1737, C₂₆H₂₄N₂O₄ requires [M]⁺ 428.1736.

General procedure for synthesis of (12a-f)

A mixture of the tetraformyl compound **11** (0.117 mmol) and the hydrazine or aryl hydrazine (0.23 mmol) in absolute ethanol (5 mL) was heated with stirring at reflux for 6-8 h. After cooling and concentrating the solution, the resulting crystals were collected by filtration and recrystallized from EtOH to give the corresponding pyrazoles.

3,3',3',3'-Tetramethyl-2,2'-di(1*H*-pyrazol-4-yl)-3*H*,3'*H*-5,5'-biindole (12a). (0.044 g, Yield 90%); mp 218-220 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 3168, 2964, 1612, 1567, 1535, 1458; ¹H NMR (DMSO-*d*₆): δ 1.50 (s, 12H), 7.53 (d, *J* = 7.8 Hz, 2H), 7.62 (d, *J* = 7.8 Hz, 2H), 7.81 (s, 2H), 8.35 (s, 4H), 12.75 (bs, 2H); ¹³C NMR (DMSO-*d*₆): δ 24.71, 53.24, 115.28, 119.81, 120.41, 126.68, 127.70, 137.73, 147.78, 153.39, 179.45; MS (EI, 70 ev): *m/z* (%) 420 (M⁺, 63), 405 (24), 370 (17), 354 (13), 302 (100), 287 (47), 44 (18). Found: [M]⁺ 420.2062, C₂₆H₂₄N₆ requires [M]⁺ 420.2062.

2,2'-Bis[1-(2-chlorophenyl)-1*H*-pyrazol-4-yl]-3,3',3',3'-tetramethyl-3*H*,3'*H*-5,5'-biindole (12b). (0.067 g, Yield 89%); mp 155-157 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2967, 2927, 1612, 1576, 1527, 1492, 1459; ¹H NMR (DMSO-*d*₆): δ 1.59 (s, 12H), 7.56-7.76 (m, 12H), 7.91 (s, 2H), 8.51 (s,

2H), 9.04 (s, 2H); ^{13}C NMR (DMSO- d_6): δ 24.58, 53.44, 116.48, 119.84, 120.75, 126.99, 128.25, 128.77, 128.83, 128.87, 131.01, 133.82, 137.71, 138.26, 141.32, 147.60, 151.97, 178.78; MS (EI, 70 eV): m/z (%) 644 (1), 642 (6), 640 (M^+ , 10), 480 (28), 467 (29), 440 (26), 412 (100), 397 (32), 111 (10). Found: $[\text{M}]^+$ 640.1909, $\text{C}_{38}\text{H}_{30}\text{Cl}_2\text{N}_6$ requires $[\text{M}]^+$ 640.1909.

2,2'-Bis[1-(3-chlorophenyl)-1H-pyrazol-4-yl]-3,3,3',3'-tetramethyl-3H,3'H-5,5'-biindole (12c). (0.064 g, Yield 85%); mp 153-155 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2967, 2928, 1620, 1593, 1529, 1487, 1461, 1408; ^1H NMR (DMSO- d_6): δ 1.61 (s, 12H), 7.43 (d, $J = 7.8$ Hz, 2H), 7.57 (d, $J = 7.8$ Hz, 2H), 7.61 (t, $J = 7.8$ Hz, 2H), 7.70 (d, $J = 7.8$ Hz, 2H), 7.91 (s, 2H), 8.03 (d, $J = 7.8$ Hz, 2H), 8.2 (s, 2H), 8.48 (s, 2H), 9.33 (s, 2H); ^{13}C NMR (DMSO- d_6): δ 24.36, 53.46, 117.95, 118.31, 119.02, 120.23, 120.59, 126.88, 127.14, 128.76, 131.74, 134.56, 138.18, 140.75, 142, 148.05, 153.02, 178.52; MS (EI, 70 eV): m/z (%) 644 (1), 642 (6), 640 (M^+ , 10), 480 (35), 467 (30), 440 (25), 412 (100), 397 (30), 111 (18). Found: $[\text{M}]^+$ 640.1908, $\text{C}_{38}\text{H}_{30}\text{Cl}_2\text{N}_6$ requires $[\text{M}]^+$ 640.1909.

2,2'-Bis[1-(4-chlorophenyl)-1H-pyrazol-4-yl]-3,3,3',3'-tetramethyl-3H,3'H-5,5'-biindole (12d). (0.071 g, Yield 95%); mp 222-225 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2968, 2927, 1610, 1577, 1529, 1499, 1463; ^1H NMR (DMSO- d_6): δ 1.14 (s, 12H), 7.54-7.61 (m, 8H), 7.84 (s, 2H), 8.01 (d, $J = 8.7$ Hz, 4H), 8.40 (s, 2H), 9.21 (s, 2H); ^{13}C NMR (DMSO- d_6): δ 24.38, 53.44, 118.27, 120.21, 120.57, 121.05, 126.85, 128.57, 129.94, 131.57, 138.13, 138.47, 141.79, 148.04, 153.13, 178.56; MS (EI, 70 eV): m/z (%) 644 (1), 642 (6), 640 (M^+ , 10), 480 (28), 467 (43), 440 (27), 412 (100), 397 (44), 111 (13). Found: $[\text{M}]^+$ 640.1909, $\text{C}_{38}\text{H}_{30}\text{Cl}_2\text{N}_6$ requires $[\text{M}]^+$ 640.1909.

2,2'-Bis[1-(4-methoxyphenyl)-1H-pyrazol-4-yl]-3,3,3',3'-tetramethyl-3H,3'H-5,5'-biindole (12e). (0.067 g, Yield 90%); mp 182-186 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2963, 2928, 1610, 1560, 1514, 1459, 1250, 1172, 1034; ^1H NMR (DMSO- d_6): δ 1.64 (s, 12H), 3.82 (s, 6H), 7.11 (d, $J = 8.4$ Hz, 4H), 7.63 (d, $J = 7.8$ Hz, 2H), 7.73 (d, $J = 7.8$ Hz, 2H), 7.88 (s, 2H), 7.92 (d, $J = 8.4$ Hz, 4H), 8.39 (s, 2H), 9.12 (s, 2H); ^{13}C NMR (DMSO- d_6): δ 24.63, 53.41, 55.98, 115.11, 116.54, 119.39, 120.86, 121.21, 127.09, 129.17, 133.05, 138.31, 141.51, 147.35, 150.85, 158.84, 178.99; MS (EI, 70 eV): m/z (%) 632 (M^+ , 2), 476 (50), 463 (58), 436 (31), 408 (100), 393 (87), 204 (33), 77 (22), 57 (23), 43 (23). Found: $[\text{M}]^+$ 632.2898, $\text{C}_{40}\text{H}_{36}\text{N}_6\text{O}_2$ requires $[\text{M}]^+$ 632.2899.

2,2'-Bis[1-(4-bromophenyl)-1H-pyrazol-4-yl]-3,3,3',3'-tetramethyl-3H,3'H-5,5'-biindole (12f). (0.074 g, Yield 87%); mp 230-233 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2969, 2927, 1611, 1577, 1529, 1495, 1463; ^1H NMR (DMSO- d_6): δ 1.63 (s, 12H), 7.64 (d, $J = 7.8$ Hz, 2H), 7.72 (d, $J = 7.8$ Hz, 2H), 7.76 (d, $J = 8.1$ Hz, 4H), 7.96 (s, 2H), 8.02 (d, $J = 8.1$ Hz, 4H), 8.57 (s, 2H), 9.37 (s, 2H); ^{13}C NMR (DMSO- d_6): δ 24.47, 53.46, 117.48, 119.73, 120.09, 120.81, 121.48, 127.06, 129.27, 132.90, 138.35, 138.74, 142.135, 147.60, 151.44, 178.71; MS (EI, 70 eV): m/z (%) 732 (2), 730 (4), 728 (M^+ , 2), 485 (54), 417 (54), 327 (23), 171 (100), 92 (43), 65 (57), 43 (35). Found: $[\text{M}]^+$ 728.0897, $[\text{M}+2]^+$ 730.0878, $\text{C}_{38}\text{H}_{30}\text{Br}_2\text{N}_6$ requires $[\text{M}]^+$ 728.0899, $[\text{M}+2]^+$ 730.0878.

2,2'-(1,1,6,6-Tetramethylindolo[7,6-g]indole-2,7(1H,3H,6H,8H)-diylidene)dimalonaldehyde (15). To *N,N*-dimethylformamide (3 mL) cooled in an ice bath was added dropwise phosphorus oxychloride (2.2 mL, 24 mmol) with stirring at below 5 °C. After this addition, a solution of **14** (1.16 g, 4 mmol) in DMF (2.5 mL) was added dropwise. The cooling bath was removed and the

reaction mixture was stirred at 75 °C for 12 h. The resulting solution was added to ice-cooled water and made alkaline with NaOH (aq.) solution. The resulting precipitate was collected by filtration after 24 h, dried in air, recrystallized from ethanol, to give **15**. (1.37 g, Yield 85%); mp 215-218 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 3130, 2924, 2854, 2766, 1679, 1614, 1592, 1516, 1308; ^1H NMR (CDCl_3): δ 1.86 (s, 12H), 7.63 (d, $J = 8.4$ Hz, 2H), 7.88 (d, $J = 8.4$ Hz, 2H), 9.83 (s, 2H), 9.87 (s, 2H), 14.35 (bs, 2H); ^{13}C NMR (CDCl_3): δ 22.87, 52.66, 110.20, 118.83, 120.50, 120.75, 135.48, 137.45, 179.94, 187.65, 192.73; MS (EI, 70 ev): m/z (%) 402 (M^+ , 68), 374 (43), 359 (80), 335 (32), 265 (57), 237 (46), 222 (100), 194 (39), 165 (30), 69 (36), 44 (98). Found: $[\text{M}]^+$ 402.1580, $\text{C}_{24}\text{H}_{22}\text{N}_2\text{O}_4$ requires $[\text{M}]^+$ 402.1579.

General procedure for synthesis of (16a-e)

A mixture of the tetraformyl compound **15** (0.2 mmol) and the hydrazine or aryl hydrazines (0.4 mmol) in absolute ethanol (10 mL) was heated with stirring at reflux for 10-12 h. After cooling and concentrating the solution, the resulting crystals were collected by filtration and recrystallized from EtOH to give the corresponding pyrazoles.

1,1,6,6-Tetramethyl-2,7-di(1H-pyrazol-4-yl)-1,6-dihydroindolo[7,6-g]indole (16a). (0.063 g, Yield 80%); mp 220-221 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 3191, 2961, 2925, 2861, 1627, 1572, 1517; ^1H NMR ($\text{DMSO}-d_6$): δ 1.52 (s, 12H), 7.74 (d, $J = 8.1$ Hz, 2H), 8.38 (d, $J = 8.1$ Hz, 2H), 8.50 (s, 4H), 13.38 (bs, 2H); ^{13}C NMR ($\text{DMSO}-d_6$): δ 24.36, 54.37, 115.56, 120.02, 120.32, 127.01, 130.95, 142.56, 149.62, 179.68; MS (EI, 70 ev): m/z (%) 394 (M^+ , 100), 379 (46), 261 (32), 83 (23), 57 (42), 43 (66). Found: $[\text{M}]^+$ 394.1906, $\text{C}_{24}\text{H}_{22}\text{N}_6$ requires $[\text{M}]^+$ 394.1906.

2,7-Bis(1-(3-chlorophenyl)-1H-pyrazol-4-yl)-1,1,6,6-tetramethyl-1,6-dihydroindolo[7,6-g]indole (16b). (0.11 g, Yield 89%); mp 234-237 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2960, 2926, 2863, 1610, 1595, 1571, 1488, 1457; ^1H NMR ($\text{DMSO}-d_6$): δ 1.60 (s, 12H), 7.42 (d, $J = 7.8$ Hz, 2H), 7.57 (t, $J = 8.1$ Hz, 2H), 7.79 (d, $J = 8.1$ Hz, 2H), 8.03 (d, $J = 7.8$ Hz, 2H), 8.20 (s, 2H), 8.44 (d, $J = 8.1$ Hz, 2H), 8.54 (s, 2H), 9.33 (s, 2H); ^{13}C NMR ($\text{DMSO}-d_6$): δ 23.97, 54.58, 117.90, 118.60, 118.98, 120.23, 120.87, 127.07, 127.11, 128.45, 131.70, 134.54, 140.78, 141.89, 143.17, 149.33, 178.64; MS (EI, 70 ev): m/z (%) 618 (1), 616 (4), 614 (M^+ , 7), 482 (18), 441 (22), 394 (24), 371 (100), 356 (31), 152 (35), 127 (40), 57 (47), 43 (67). Found: $[\text{M}]^+$ 614.1754, $\text{C}_{36}\text{H}_{28}\text{Cl}_2\text{N}_6$ requires $[\text{M}]^+$ 614.1753.

2,7-Bis(1-(4-chlorophenyl)-1H-pyrazol-4-yl)-1,1,6,6-tetramethyl-1,6-dihydroindolo[7,6-g]indole (16c). (0.108 g, Yield 88%); mp 213-215 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2964, 2926, 2863, 1612, 1567, 1495, 949; ^1H NMR ($\text{DMSO}-d_6$): δ 1.63 (s, 12H), 7.64 (d, $J = 8.4$ Hz, 4H), 7.86 (d, $J = 8.1$ Hz, 2H), 8.10 (d, $J = 8.4$ Hz, 4H), 8.54 (d, $J = 8.1$ Hz, 2H), 8.64 (s, 2H), 9.38 (s, 2H); ^{13}C NMR ($\text{DMSO}-d_6$): δ 24.15, 54.57, 117.37, 120.63, 121.25, 121.50, 126.06, 129.54, 129.72, 130, 131.88, 138.31, 142.27, 143.08, 179.30; MS (EI, 70 ev): m/z (%) 618 (1), 616 (4), 614 (M^+ , 7), 394 (5), 371 (4), 279 (8), 167 (17), 149 (51), 127 (100), 84 (36), 71 (43), 57 (75), 43 (90). Found: $[\text{M}]^+$ 614.1751, $\text{C}_{36}\text{H}_{28}\text{Cl}_2\text{N}_6$ requires $[\text{M}]^+$ 614.1753.

2,7-Bis(1-(4-methoxyphenyl)-1H-pyrazol-4-yl)-1,1,6,6-tetramethyl-1,6-dihydroindolo[7,6-g]indole (16d). (0.115 g, Yield 95%); mp 223-225 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2971, 2930, 1612,

1578, 1515, 1255, 1175; ^1H NMR (DMSO- d_6): δ 1.64 (s, 12H), 3.82 (s, 6H), 7.12 (d, J = 8.4 Hz, 4H), 7.88 (d, J = 8.7 Hz, 2H), 7.95 (d, J = 8.4 Hz, 4H), 8.60 (d, J = 8.7 Hz, 2H), 8.65 (s, 2H), 9.33 (s, 2H); ^{13}C NMR (DMSO- d_6): δ 24.26, 54.51, 55.98, 115.11, 116.16, 116.82, 120.55, 121.21, 122.24, 126.24, 129.04, 133.09, 141.52, 142.91, 158.81, 179.45; MS (EI, 70 eV): m/z (%) 606 (M^+ , 100), 591 (30), 434 (9), 367 (16), 303 (20), 108 (16), 77 (14), 44 (16). Found: $[\text{M}]^+$ 606.2744, $\text{C}_{38}\text{H}_{34}\text{N}_6\text{O}_2$ requires $[\text{M}]^+$ 606.2743.

2,7-Bis(1-(4-bromophenyl)-1H-pyrazol-4-yl)-1,1,6,6-tetramethyl-1,6-dihydroindolo[7,6-g]indole (16e). (0.122 g, Yield 87%); mp 229-230 °C; FT-IR (KBr) $\nu_{\text{max}}/\text{cm}^{-1}$: 2969, 2927, 1619, 1573, 1524, 1494; ^1H NMR (DMSO- d_6): δ 1.62 (s, 12H), 7.77 (d, J = 8.4 Hz, 4H), 7.85 (d, J = 8.1 Hz, 2H), 8.05 (d, J = 8.4 Hz, 4H), 8.51 (d, J = 8.1 Hz, 2H), 8.61 (s, 2H), 9.36 (s, 2H); ^{13}C NMR (DMSO- d_6): δ 24.10, 54.58, 120.06, 120.54, 121.48, 126.48, 129.16, 132.21, 132.44, 132.63, 132.90, 138.77, 142.14, 143.12, 179.12; MS (EI, 70 eV): m/z (%) 706 (4), 704 (8), 702 (M^+ , 4), 614 (20), 524 (34), 484 (32), 458 (100), 441 (40), 229 (18), 171 (25), 65 (23), 43 (41). Found: $[\text{M}]^+$ 702.0744, $[\text{M}+2]^+$ 704.0721, $\text{C}_{36}\text{H}_{28}\text{Br}_2\text{N}_6$ requires $[\text{M}]^+$ 702.0742, $[\text{M}+2]^+$ 704.0722.

Acknowledgements

The authors are grateful to the University of Urmia for financial support of this work. We also thank Mr. M. Qavidel for measuring the NMR spectra.

References

1. Baradarani, M. M.; Afghan, A.; Zebarjadi, F.; Hasanzadeh, K.; Joule, J. A. *J. Heterocycl. Chem.* **2006**, *43*, 1591.
2. Helliwell, M.; Afgan, A.; Baradarani, M. M.; Joule, J. A. *Acta Crystallogr. Sect. E* **2006**, *62*, o737.
3. Rashidi, A.; Afghan, A.; Baradarani, M. M.; Joule, J. A. *J. Heterocycl. Chem.* **2009**, *46*, 428.
4. Helliwell, M.; Afghan, A.; Keshvari, F.; Baradarani, M. M.; Joule, J. A. *Acta Crystallogr. Sect. E* **2010**, *66*, o112.
5. Ferlin, M. G.; Chiarello, G.; Basadonna, O.; Gia, O.; Mobilo, S.; Baccichetti, F.; Carlassare, F. *Farmaco* **1989**, *44*, 1141.
6. Robinson, B. *The Fischer Indole Synthesis*; Wiley: New York, 1982.
7. Kolesnikov, A. M.; Mikhailenko, F. A. *Zh. Org. Khim.* **1982**, *18*, 441.
8. Samsoniya, Sh. A.; Trapaidze, M. V.; Machaidze, N. N.; Dürr, G. *Chem. Heterocycl. Comp.* **2002**, *38*, 4.